

# Appendix A Review of Fundamental Concepts of Algebra

## A.1 Real Numbers and Their Properties

### What you should learn

- Represent and classify real numbers.
- Order real numbers and use inequalities.
- Find the absolute values of real numbers and find the distance between two real numbers.
- Evaluate algebraic expressions.
- Use the basic rules and properties of algebra.

### Why you should learn it

Real numbers are used to represent many real-life quantities. For example, in Exercise 65 on page A9, you will use real numbers to represent the federal deficit.

The *HM mathSpace*® CD-ROM and *Eduspace*® for this text contain additional resources related to the concepts discussed in this chapter.




FIGURE A.1 Subsets of real numbers

### Real Numbers

**Real numbers** are used in everyday life to describe quantities such as age, miles per gallon, and population. Real numbers are represented by symbols such as

$$-5, 9, 0, \frac{4}{3}, 0.666 \dots, 28.21, \sqrt{2}, \pi, \text{ and } \sqrt[3]{-32}.$$

Here are some important **subsets** (each member of subset  $B$  is also a member of set  $A$ ) of the real numbers. The three dots, called *ellipsis points*, indicate that the pattern continues indefinitely.

$$\{1, 2, 3, 4, \dots\} \quad \text{Set of natural numbers}$$

$$\{0, 1, 2, 3, 4, \dots\} \quad \text{Set of whole numbers}$$

$$\{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\} \quad \text{Set of integers}$$

A real number is **rational** if it can be written as the ratio  $p/q$  of two integers, where  $q \neq 0$ . For instance, the numbers

$$\frac{1}{3} = 0.3333 \dots = 0.\overline{3}, \frac{1}{8} = 0.125, \text{ and } \frac{125}{111} = 1.126126 \dots = 1.\overline{126}$$

are rational. The decimal representation of a rational number either repeats (as in  $\frac{173}{55} = 3.14\overline{5}$ ) or terminates (as in  $\frac{1}{2} = 0.5$ ). A real number that cannot be written as the ratio of two integers is called **irrational**. Irrational numbers have infinite nonrepeating decimal representations. For instance, the numbers

$$\sqrt{2} = 1.4142135 \dots \approx 1.41 \quad \text{and} \quad \pi = 3.1415926 \dots \approx 3.14$$

are irrational. (The symbol  $\approx$  means “is approximately equal to.”) Figure A.1 shows subsets of real numbers and their relationships to each other.

Real numbers are represented graphically by a **real number line**. The point 0 on the real number line is the **origin**. Numbers to the right of 0 are positive, and numbers to the left of 0 are negative, as shown in Figure A.2. The term **nonnegative** describes a number that is either positive or zero.





FIGURE A.2 The real number line

As illustrated in Figure A.3, there is a *one-to-one correspondence* between real numbers and points on the real number line.



Every real number corresponds to exactly one point on the real number line.

FIGURE A.3 One-to-one

Every point on the real number line corresponds to exactly one real number.

## Ordering Real Numbers

One important property of real numbers is that they are *ordered*.

### Definition of Order on the Real Number Line

If  $a$  and  $b$  are real numbers,  $a$  is less than  $b$  if  $b - a$  is positive. The **order** of  $a$  and  $b$  is denoted by the **inequality**  $a < b$ . This relationship can also be described by saying that  $b$  is *greater than*  $a$  and writing  $b > a$ . The inequality  $a \leq b$  means that  $a$  is *less than or equal to*  $b$ , and the inequality  $b \geq a$  means that  $b$  is *greater than or equal to*  $a$ . The symbols  $<$ ,  $>$ ,  $\leq$ , and  $\geq$  are *inequality symbols*.




FIGURE A.4  $a < b$  if and only if  $a$  lies to the left of  $b$ .




FIGURE A.5




FIGURE A.6

Geometrically, this definition implies that  $a < b$  if and only if  $a$  lies to the left of  $b$  on the real number line, as shown in Figure A.4.

### Example 1 Interpreting Inequalities

Describe the subset of real numbers represented by each inequality.

- a.  $x \leq 2$       b.  $-2 \leq x < 3$

#### Solution

- a. The inequality  $x \leq 2$  denotes all real numbers less than or equal to 2, as shown in Figure A.5.  
 b. The inequality  $-2 \leq x < 3$  means that  $x \geq -2$  and  $x < 3$ . This “double inequality” denotes all real numbers between  $-2$  and  $3$ , including  $-2$  but not including  $3$ , as shown in Figure A.6.

**CHECKPOINT** Now try Exercise 19.

Inequalities can be used to describe subsets of real numbers called **intervals**. In the bounded intervals below, the real numbers  $a$  and  $b$  are the **endpoints** of each interval. The endpoints of a closed interval are included in the interval, whereas the endpoints of an open interval are not included in the interval.

### STUDY TIP

The reason that the four types of intervals at the right are called *bounded* is that each has a finite length. An interval that does not have a finite length is *unbounded* (see page A3).

### Bounded Intervals on the Real Number Line






Notation	Interval Type	Inequality	Graph
$[a, b]$	Closed	$a \leq x \leq b$	
$(a, b)$	Open	$a < x < b$	
$[a, b)$		$a \leq x < b$	
$(a, b]$		$a < x \leq b$	

## STUDY TIP

Note that whenever you write intervals containing  $\infty$  or  $-\infty$ , you always use a parenthesis and never a bracket. This is because these symbols are never an endpoint of an interval and therefore not included in the interval.

The symbols  $\infty$ , **positive infinity**, and  $-\infty$ , **negative infinity**, do not represent real numbers. They are simply convenient symbols used to describe the unboundedness of an interval such as  $(1, \infty)$  or  $(-\infty, 3]$ .

### Unbounded Intervals on the Real Number Line

Notation	Interval Type	Inequality	Graph
$[a, \infty)$		$x \geq a$	
$(a, \infty)$	Open	$x > a$	
$(-\infty, b]$		$x \leq b$	
$(-\infty, b)$	Open	$x < b$	
$(-\infty, \infty)$	Entire real line	$-\infty < x < \infty$	

### Example 2 Using Inequalities to Represent Intervals

Use inequality notation to describe each of the following.

- a.  $c$  is at most 2.      b.  $m$  is at least  $-3$ .  
 c. All  $x$  in the interval  $(-3, 5]$

#### Solution

- a. The statement “ $c$  is at most 2” can be represented by  $c \leq 2$ .  
 b. The statement “ $m$  is at least  $-3$ ” can be represented by  $m \geq -3$ .  
 c. “All  $x$  in the interval  $(-3, 5]$ ” can be represented by  $-3 < x \leq 5$ .

 **CHECKPOINT** Now try Exercise 31.

### Example 3 Interpreting Intervals

Give a verbal description of each interval.

- a.  $(-1, 0)$       b.  $[2, \infty)$       c.  $(-\infty, 0)$

#### Solution

- a. This interval consists of all real numbers that are greater than  $-1$  and less than  $0$ .  
 b. This interval consists of all real numbers that are greater than or equal to  $2$ .  
 c. This interval consists of all negative real numbers.

 **CHECKPOINT** Now try Exercise 29.

The **Law of Trichotomy** states that for any two real numbers  $a$  and  $b$ , precisely one of three relationships is possible:

$$a = b, \quad a < b, \quad \text{or} \quad a > b. \quad \text{Law of Trichotomy}$$

## Absolute Value and Distance

The **absolute value** of a real number is its *magnitude*, or the distance between the origin and the point representing the real number on the real number line.

### Definition of Absolute Value

If  $a$  is a real number, then the absolute value of  $a$  is

$$|a| = \begin{cases} a, & \text{if } a \geq 0 \\ -a, & \text{if } a < 0 \end{cases}$$

Notice in this definition that the absolute value of a real number is never negative. For instance, if  $a = -5$ , then  $|-5| = -(-5) = 5$ . The absolute value of a real number is either positive or zero. Moreover, 0 is the only real number whose absolute value is 0. So,  $|0| = 0$ .

### Example 4 Evaluating the Absolute Value of a Number

Evaluate  $\frac{|x|}{x}$  for (a)  $x > 0$  and (b)  $x < 0$ .

#### Solution

a. If  $x > 0$ , then  $|x| = x$  and  $\frac{|x|}{x} = \frac{x}{x} = 1$ .

b. If  $x < 0$ , then  $|x| = -x$  and  $\frac{|x|}{x} = \frac{-x}{x} = -1$ .

 **CHECKPOINT** Now try Exercise 47.

### Properties of Absolute Values

1.  $|a| \geq 0$
2.  $|-a| = |a|$
3.  $|ab| = |a||b|$
4.  $\left|\frac{a}{b}\right| = \frac{|a|}{|b|}$ ,  $b \neq 0$

Absolute value can be used to define the distance between two points on the real number line. For instance, the distance between  $-3$  and  $4$  is

$$\begin{aligned} |-3 - 4| &= |-7| \\ &= 7 \end{aligned}$$

as shown in Figure A.7.



FIGURE A.7 The distance between  $-3$  and  $4$  is  $7$ .

### Distance Between Two Points on the Real Number Line

Let  $a$  and  $b$  be real numbers. The **distance between  $a$  and  $b$**  is

$$d(a, b) = |b - a| = |a - b|.$$

## Algebraic Expressions

One characteristic of algebra is the use of letters to represent numbers. The letters are **variables**, and combinations of letters and numbers are **algebraic expressions**. Here are a few examples of algebraic expressions.

$$5x, \quad 2x - 3, \quad \frac{4}{x^2 + 2}, \quad 7x + y$$

### Definition of an Algebraic Expression

An **algebraic expression** is a collection of letters (**variables**) and real numbers (**constants**) combined using the operations of addition, subtraction, multiplication, division, and exponentiation.

The **terms** of an algebraic expression are those parts that are separated by *addition*. For example,

$$x^2 - 5x + 8 = x^2 + (-5x) + 8$$

has three terms:  $x^2$  and  $-5x$  are the **variable terms** and 8 is the **constant term**. The numerical factor of a variable term is the **coefficient** of the variable term. For instance, the coefficient of  $-5x$  is  $-5$ , and the coefficient of  $x^2$  is 1.

To **evaluate** an algebraic expression, substitute numerical values for each of the variables in the expression. Here are two examples.

<i>Expression</i>	<i>Value of Variable</i>	<i>Substitute</i>	<i>Value of Expression</i>
$-3x + 5$	$x = 3$	$-3(3) + 5$	$-9 + 5 = -4$
$3x^2 + 2x - 1$	$x = -1$	$3(-1)^2 + 2(-1) - 1$	$3 - 2 - 1 = 0$

When an algebraic expression is evaluated, the **Substitution Principle** is used. It states that “If  $a = b$ , then  $a$  can be replaced by  $b$  in any expression involving  $a$ .” In the first evaluation shown above, for instance, 3 is *substituted* for  $x$  in the expression  $-3x + 5$ .

## Basic Rules of Algebra

There are four arithmetic operations with real numbers: *addition*, *multiplication*, *subtraction*, and *division*, denoted by the symbols  $+$ ,  $\times$  or  $\cdot$ ,  $-$ , and  $\div$  or  $/$ . Of these, addition and multiplication are the two primary operations. Subtraction and division are the inverse operations of addition and multiplication, respectively.

### Definitions of Subtraction and Division

**Subtraction:** Add the opposite.

**Division:** Multiply by the reciprocal.

$$a - b = a + (-b)$$

$$\text{If } b \neq 0, \text{ then } a/b = a\left(\frac{1}{b}\right) = \frac{a}{b}.$$

In these definitions,  $-b$  is the **additive inverse** (or opposite) of  $b$ , and  $1/b$  is the **multiplicative inverse** (or reciprocal) of  $b$ . In the fractional form  $a/b$ ,  $a$  is the **numerator** of the fraction and  $b$  is the **denominator**.

Because the properties of real numbers below are true for variables and algebraic expressions as well as for real numbers, they are often called the **Basic Rules of Algebra**. Try to formulate a verbal description of each property. For instance, the first property states that *the order in which two real numbers are added does not affect their sum*.

### Basic Rules of Algebra

Let  $a$ ,  $b$ , and  $c$  be real numbers, variables, or algebraic expressions.

	<i>Property</i>	<i>Example</i>
Commutative Property of Addition:	$a + b = b + a$	$4x + x^2 = x^2 + 4x$
Commutative Property of Multiplication:	$ab = ba$	$(4 - x)x^2 = x^2(4 - x)$
Associative Property of Addition:	$(a + b) + c = a + (b + c)$	$(x + 5) + x^2 = x + (5 + x^2)$
Associative Property of Multiplication:	$(ab)c = a(bc)$	$(2x \cdot 3y)(8) = (2x)(3y \cdot 8)$
Distributive Properties:	$a(b + c) = ab + ac$	$3x(5 + 2x) = 3x \cdot 5 + 3x \cdot 2x$
	$(a + b)c = ac + bc$	$(y + 8)y = y \cdot y + 8 \cdot y$
Additive Identity Property:	$a + 0 = a$	$5y^2 + 0 = 5y^2$
Multiplicative Identity Property:	$a \cdot 1 = a$	$(4x^2)(1) = 4x^2$
Additive Inverse Property:	$a + (-a) = 0$	$5x^3 + (-5x^3) = 0$
Multiplicative Inverse Property:	$a \cdot \frac{1}{a} = 1, \quad a \neq 0$	$(x^2 + 4) \left( \frac{1}{x^2 + 4} \right) = 1$

Because subtraction is defined as “adding the opposite,” the Distributive Properties are also true for subtraction. For instance, the “subtraction form” of  $a(b + c) = ab + ac$  is  $a(b - c) = ab - ac$ .

### STUDY TIP

Notice the difference between the *opposite of a number* and a *negative number*. If  $a$  is already negative, then its opposite,  $-a$ , is positive. For instance, if  $a = -5$ , then

$$-a = -(-5) = 5.$$

### Properties of Negation and Equality

Let  $a$  and  $b$  be real numbers, variables, or algebraic expressions.

	<i>Property</i>	<i>Example</i>
1.	$(-1)a = -a$	$(-1)7 = -7$
2.	$-(-a) = a$	$-(-6) = 6$
3.	$(-a)b = -(ab) = a(-b)$	$(-5)3 = -(5 \cdot 3) = 5(-3)$
4.	$(-a)(-b) = ab$	$(-2)(-x) = 2x$
5.	$-(a + b) = (-a) + (-b)$	$-(x + 8) = (-x) + (-8)$ $= -x - 8$
6.	If $a = b$ , then $a \pm c = b \pm c$ .	$\frac{1}{2} + 3 = 0.5 + 3$
7.	If $a = b$ , then $ac = bc$ .	$4^2 \cdot 2 = 16 \cdot 2$
8.	If $a \pm c = b \pm c$ , then $a = b$ .	$1.4 - 1 = \frac{7}{5} - 1 \Rightarrow 1.4 = \frac{7}{5}$
9.	If $ac = bc$ and $c \neq 0$ , then $a = b$ .	$3x = 3 \cdot 4 \Rightarrow x = 4$

## STUDY TIP

The “or” in the Zero-Factor Property includes the possibility that either or both factors may be zero. This is an **inclusive or**, and it is the way the word “or” is generally used in mathematics.

## Properties of Zero

Let  $a$  and  $b$  be real numbers, variables, or algebraic expressions.

- $a + 0 = a$  and  $a - 0 = a$
- $a \cdot 0 = 0$
- $\frac{0}{a} = 0$ ,  $a \neq 0$
- $\frac{a}{0}$  is undefined.
- Zero-Factor Property:** If  $ab = 0$ , then  $a = 0$  or  $b = 0$ .

## Properties and Operations of Fractions

Let  $a$ ,  $b$ ,  $c$ , and  $d$  be real numbers, variables, or algebraic expressions such that  $b \neq 0$  and  $d \neq 0$ .

- Equivalent Fractions:**  $\frac{a}{b} = \frac{c}{d}$  if and only if  $ad = bc$ .
- Rules of Signs:**  $-\frac{a}{b} = \frac{-a}{b} = \frac{a}{-b}$  and  $\frac{-a}{-b} = \frac{a}{b}$
- Generate Equivalent Fractions:**  $\frac{a}{b} = \frac{ac}{bc}$ ,  $c \neq 0$
- Add or Subtract with Like Denominators:**  $\frac{a}{b} \pm \frac{c}{b} = \frac{a \pm c}{b}$
- Add or Subtract with Unlike Denominators:**  $\frac{a}{b} \pm \frac{c}{d} = \frac{ad \pm bc}{bd}$
- Multiply Fractions:**  $\frac{a}{b} \cdot \frac{c}{d} = \frac{ac}{bd}$
- Divide Fractions:**  $\frac{a}{b} \div \frac{c}{d} = \frac{a}{b} \cdot \frac{d}{c} = \frac{ad}{bc}$ ,  $c \neq 0$

## STUDY TIP

In Property 1 of fractions, the phrase “if and only if” implies two statements. One statement is: If  $a/b = c/d$ , then  $ad = bc$ . The other statement is: If  $ad = bc$ , where  $b \neq 0$  and  $d \neq 0$ , then  $a/b = c/d$ .

### Example 5 Properties and Operations of Fractions

- a. Equivalent fractions:  $\frac{x}{5} = \frac{3 \cdot x}{3 \cdot 5} = \frac{3x}{15}$     b. Divide fractions:  $\frac{7}{x} \div \frac{3}{2} = \frac{7}{x} \cdot \frac{2}{3} = \frac{14}{3x}$
- c. Add fractions with unlike denominators:  $\frac{x}{3} + \frac{2x}{5} = \frac{5 \cdot x + 3 \cdot 2x}{3 \cdot 5} = \frac{11x}{15}$

 **CHECKPOINT** Now try Exercise 103.

If  $a$ ,  $b$ , and  $c$  are integers such that  $ab = c$ , then  $a$  and  $b$  are **factors** or **divisors** of  $c$ . A **prime number** is an integer that has exactly two positive factors—itsself and 1—such as 2, 3, 5, 7, and 11. The numbers 4, 6, 8, 9, and 10 are **composite** because each can be written as the product of two or more prime numbers. The number 1 is neither prime nor composite. The **Fundamental Theorem of Arithmetic** states that every positive integer greater than 1 can be written as the product of prime numbers in precisely one way (disregarding order). For instance, the *prime factorization* of 24 is  $24 = 2 \cdot 2 \cdot 2 \cdot 3$ .

**A.1 Exercises** The *HM mathSpace*® CD-ROM and *Eduspace*® for this text contain step-by-step solutions to all odd-numbered exercises. They also provide Tutorial Exercises for additional help.

**VOCABULARY CHECK:** Fill in the blanks.

1. A real number is \_\_\_\_\_ if it can be written as the ratio  $\frac{p}{q}$  of two integers, where  $q \neq 0$ .
2. \_\_\_\_\_ numbers have infinite nonrepeating decimal representations.
3. The distance between a point on the real number line and the origin is the \_\_\_\_\_ of the real number.
4. A number that can be written as the product of two or more prime numbers is called a \_\_\_\_\_ number.
5. An integer that has exactly two positive factors, the integer itself and 1, is called a \_\_\_\_\_ number.
6. An algebraic expression is a collection of letters called \_\_\_\_\_ and real numbers called \_\_\_\_\_.
7. The \_\_\_\_\_ of an algebraic expression are those parts separated by addition.
8. The numerical factor of a variable term is the \_\_\_\_\_ of the variable term.
9. The \_\_\_\_\_ states that if  $ab = 0$ , then  $a = 0$  or  $b = 0$ .


In Exercises 1–6, determine which numbers in the set are (a) natural numbers, (b) whole numbers, (c) integers, (d) rational numbers, and (e) irrational numbers.

1.  $-9, -\frac{7}{2}, 5, \frac{2}{3}, \sqrt{2}, 0, 1, -4, 2, -11$
2.  $\sqrt{5}, -7, -\frac{7}{3}, 0, 3.12, \frac{5}{4}, -3, 12, 5$
3.  $2.01, 0.666 \dots, -13, 0.010110111 \dots, 1, -6$
4.  $2.3030030003 \dots, 0.7575, -4.63, \sqrt{10}, -75, 4$
5.  $-\pi, -\frac{1}{3}, \frac{6}{3}, \frac{1}{2}\sqrt{2}, -7.5, -1, 8, -22$
6.  $25, -17, -\frac{12}{5}, \sqrt{9}, 3.12, \frac{1}{2}\pi, 7, -11.1, 13$

In Exercises 7–10, use a calculator to find the decimal form of the rational number. If it is a nonterminating decimal, write the repeating pattern.

- |                     |                    |
|---------------------|--------------------|
| 7. $\frac{5}{8}$    | 8. $\frac{1}{3}$   |
| 9. $\frac{41}{333}$ | 10. $\frac{6}{11}$ |

In Exercises 11 and 12, approximate the numbers and place the correct symbol ( $<$  or  $>$ ) between them.



In Exercises 13–18, plot the two real numbers on the real number line. Then place the appropriate inequality symbol ( $<$  or  $>$ ) between them.

- |                                |                                  |
|--------------------------------|----------------------------------|
| 13. $-4, -8$                   | 14. $-3.5, 1$                    |
| 15. $\frac{3}{2}, 7$           | 16. $1, \frac{16}{3}$            |
| 17. $\frac{5}{6}, \frac{2}{3}$ | 18. $-\frac{8}{7}, -\frac{3}{7}$ |

In Exercises 19–30, (a) give a verbal description of the subset of real numbers represented by the inequality or the interval, (b) sketch the subset on the real number line, and (c) state whether the interval is bounded or unbounded.

- |                     |                       |
|---------------------|-----------------------|
| 19. $x \leq 5$      | 20. $x \geq -2$       |
| 21. $x < 0$         | 22. $x > 3$           |
| 23. $[4, \infty)$   | 24. $(-\infty, 2)$    |
| 25. $-2 < x < 2$    | 26. $0 \leq x \leq 5$ |
| 27. $-1 \leq x < 0$ | 28. $0 < x \leq 6$    |
| 29. $[-2, 5)$       | 30. $(-1, 2]$         |

In Exercises 31–38, use inequality notation to describe the set.

31. All  $x$  in the interval  $(-2, 4]$
32. All  $y$  in the interval  $[-6, 0)$
33.  $y$  is nonnegative.
34.  $y$  is no more than 25.
35.  $t$  is at least 10 and at most 22.
36.  $k$  is less than 5 but no less than  $-3$ .
37. The dog's weight  $W$  is more than 65 pounds.
38. The annual rate of inflation  $r$  is expected to be at least 2.5% but no more than 5%.

In Exercises 39–48, evaluate the expression.

- |                                     |                                    |
|-------------------------------------|------------------------------------|
| 39. $ -10 $                         | 40. $ 0 $                          |
| 41. $ 3 - 8 $                       | 42. $ 4 - 1 $                      |
| 43. $ -1  -  -2 $                   | 44. $-3 -  -3 $                    |
| 45. $\frac{-5}{ -5 }$               | 46. $-3 -3 $                       |
| 47. $\frac{ x + 2 }{x + 2}, x < -2$ | 48. $\frac{ x - 1 }{x - 1}, x > 1$ |

In Exercises 49–54, place the correct symbol ( $<$ ,  $>$ , or  $=$ ) between the pair of real numbers.

- 49.  $|-3|$    $-|-3|$
- 50.  $|-4|$    $|4|$
- 51.  $-5$    $-|5|$
- 52.  $-|-6|$    $|-6|$
- 53.  $-|-2|$    $-|2|$
- 54.  $-(-2)$    $-2$


In Exercises 55–60, find the distance between  $a$  and  $b$ .

- 55.  $a = 126, b = 75$
- 56.  $a = -126, b = -75$
- 57.  $a = -\frac{5}{2}, b = 0$
- 58.  $a = \frac{1}{4}, b = \frac{11}{4}$
- 59.  $a = \frac{16}{5}, b = \frac{112}{75}$
- 60.  $a = 9.34, b = -5.65$


**Budget Variance** In Exercises 61–64, the accounting department of a sports drink bottling company is checking to see whether the actual expenses of a department differ from the budgeted expenses by more than \$500 or by more than 5%. Fill in the missing parts of the table, and determine whether each actual expense passes the “budget variance test.”

	Budgeted Expense, $b$	Actual Expense, $a$	$ a - b $	$0.05b$
61. Wages	\$112,700	\$113,356	<input type="text"/>	<input type="text"/>
62. Utilities	\$9,400	\$9,772	<input type="text"/>	<input type="text"/>
63. Taxes	\$37,640	\$37,335	<input type="text"/>	<input type="text"/>
64. Insurance	\$2,575	\$2,613	<input type="text"/>	<input type="text"/>

65. **Federal Deficit** The bar graph shows the federal government receipts (in billions of dollars) for selected years from 1960 through 2000. (Source: U.S. Office of Management and Budget)




(a) Complete the table. (Hint: Find  $|\text{Receipts} - \text{Expenditures}|$ .)



Year	Expenditures (in billions)	Surplus or deficit (in billions)
1960	\$92.2	<input type="text"/>
1970	\$195.6	<input type="text"/>
1980	\$590.9	<input type="text"/>
1990	\$1253.2	<input type="text"/>
2000	\$1788.8	<input type="text"/>

(b) Use the table in part (a) to construct a bar graph showing the magnitude of the surplus or deficit for each year.

66. **Veterans** The table shows the number of living veterans (in thousands) in the United States in 2002 by age group. Construct a circle graph showing the percent of living veterans by age group as a fraction of the total number of living veterans. (Source: Department of Veteran Affairs)



Age group	Number of veterans
Under 35	2213
35–44	3290
45–54	4666
55–64	5665
65 and older	9784

In Exercises 67–72, use absolute value notation to describe the situation.

- 67. The distance between  $x$  and 5 is no more than 3.
- 68. The distance between  $x$  and  $-10$  is at least 6.
- 69.  $y$  is at least six units from 0.
- 70.  $y$  is at most two units from  $a$ .
- 71. While traveling on the Pennsylvania Turnpike, you pass milepost 326 near Valley Forge, then milepost 351 near Philadelphia. How many miles do you travel during that time period?
- 72. The temperature in Chicago, Illinois was  $48^\circ$  last night at midnight, then  $82^\circ$  at noon today. What was the change in temperature over the 12-hour period?

In Exercises 73–78, identify the terms. Then identify the coefficients of the variable terms of the expression.

73.  $7x + 4$                       74.  $6x^3 - 5x$   
 75.  $\sqrt{3}x^2 - 8x - 11$         76.  $3\sqrt{3}x^2 + 1$   
 77.  $4x^3 + \frac{x}{2} - 5$                 78.  $3x^4 - \frac{x^2}{4}$

In Exercises 79–84, evaluate the expression for each value of  $x$ . (If not possible, state the reason.)

- | <i>Expression</i>         | <i>Values</i> |              |
|---------------------------|---------------|--------------|
| 79. $4x - 6$              | (a) $x = -1$  | (b) $x = 0$  |
| 80. $9 - 7x$              | (a) $x = -3$  | (b) $x = 3$  |
| 81. $x^2 - 3x + 4$        | (a) $x = -2$  | (b) $x = 2$  |
| 82. $-x^2 + 5x - 4$       | (a) $x = -1$  | (b) $x = 1$  |
| 83. $\frac{x + 1}{x - 1}$ | (a) $x = 1$   | (b) $x = -1$ |
| 84. $\frac{x}{x + 2}$     | (a) $x = 2$   | (b) $x = -2$ |

In Exercises 85–96, identify the rule(s) of algebra illustrated by the statement.

85.  $x + 9 = 9 + x$                 86.  $2(\frac{1}{2}) = 1$   
 87.  $\frac{1}{h + 6}(h + 6) = 1, \quad h \neq -6$   
 88.  $(x + 3) - (x + 3) = 0$   
 89.  $2(x + 3) = 2 \cdot x + 2 \cdot 3$   
 90.  $(z - 2) + 0 = z - 2$   
 91.  $1 \cdot (1 + x) = 1 + x$   
 92.  $(z + 5)x = z \cdot x + 5 \cdot x$   
 93.  $x + (y + 10) = (x + y) + 10$   
 94.  $x(3y) = (x \cdot 3)y = (3x)y$   
 95.  $3(t - 4) = 3 \cdot t - 3 \cdot 4$   
 96.  $\frac{1}{7}(7 \cdot 12) = (\frac{1}{7} \cdot 7)12 = 1 \cdot 12 = 12$

In Exercises 97–104, perform the operation(s). (Write fractional answers in simplest form.)

97.  $\frac{3}{16} + \frac{5}{16}$                       98.  $\frac{6}{7} - \frac{4}{7}$   
 99.  $\frac{5}{8} - \frac{5}{12} + \frac{1}{6}$                 100.  $\frac{10}{11} + \frac{6}{33} - \frac{13}{66}$   
 101.  $12 \div \frac{1}{4}$                       102.  $-(6 \cdot \frac{4}{8})$   
 103.  $\frac{2x}{3} - \frac{x}{4}$                       104.  $\frac{5x}{6} \cdot \frac{2}{9}$

105. (a) Use a calculator to complete the table.

$n$	1	0.5	0.01	0.0001	0.000001
$5/n$					

(b) Use the result from part (a) to make a conjecture about the value of  $5/n$  as  $n$  approaches 0.

106. (a) Use a calculator to complete the table.

$n$	1	10	100	10,000	100,000
$5/n$					


(b) Use the result from part (a) to make a conjecture about the value of  $5/n$  as  $n$  increases without bound.

### Synthesis

**True or False?** In Exercises 107 and 108, determine whether the statement is true or false. Justify your answer.

107. If  $a < b$ , then  $\frac{1}{a} < \frac{1}{b}$ , where  $a \neq b \neq 0$ .  
 108. Because  $\frac{a + b}{c} = \frac{a}{c} + \frac{b}{c}$ , then  $\frac{c}{a + b} = \frac{c}{a} + \frac{c}{b}$ .  
 109. **Exploration** Consider  $|u + v|$  and  $|u| + |v|$ , where  $u \neq v \neq 0$ .  
 (a) Are the values of the expressions always equal? If not, under what conditions are they unequal?  
 (b) If the two expressions are not equal for certain values of  $u$  and  $v$ , is one of the expressions always greater than the other? Explain.  
 110. **Think About It** Is there a difference between saying that a real number is positive and saying that a real number is nonnegative? Explain.  
 111. **Think About It** Because every even number is divisible by 2, is it possible that there exist any even prime numbers? Explain.  
 112. **Writing** Describe the differences among the sets of natural numbers, whole numbers, integers, rational numbers, and irrational numbers.

In Exercises 113 and 114, use the real numbers  $A$ ,  $B$ , and  $C$  shown on the number line. Determine the sign of each expression.



113. (a)  $-A$                                       114. (a)  $-C$   
 (b)  $B - A$                                       (b)  $A - C$   
 115. **Writing** Can it ever be true that  $|a| = -a$  for a real number  $a$ ? Explain.