

10.3 POWER METHOD FOR APPROXIMATING EIGENVALUES

In Chapter 7 you saw that the eigenvalues of an $n \times n$ matrix A are obtained by solving its characteristic equation

$$\lambda^n + c_{n-1}\lambda^{n-1} + c_{n-2}\lambda^{n-2} + \cdots + c_0 = 0.$$

For large values of n , polynomial equations like this one are difficult and time-consuming to solve. Moreover, numerical techniques for approximating roots of polynomial equations of high degree are sensitive to rounding errors. In this section you will look at an alternative method for approximating eigenvalues. As presented here, the method can be used only to find the eigenvalue of A that is largest in absolute value—this eigenvalue is called the **dominant eigenvalue** of A . Although this restriction may seem severe, dominant eigenvalues are of primary interest in many physical applications.

Definition of Dominant Eigenvalue and Dominant Eigenvector

Let $\lambda_1, \lambda_2, \dots, \lambda_n$ be the eigenvalues of an $n \times n$ matrix A . λ_1 is called the **dominant eigenvalue** of A if

$$|\lambda_1| > |\lambda_i|, \quad i = 2, \dots, n.$$

The eigenvectors corresponding to λ_1 are called **dominant eigenvectors** of A .

Not every matrix has a dominant eigenvalue. For instance, the matrix

$$A = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$$

(with eigenvalues of $\lambda_1 = 1$ and $\lambda_2 = -1$) has no dominant eigenvalue. Similarly, the matrix

$$A = \begin{bmatrix} 2 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

(with eigenvalues of $\lambda_1 = 2$, $\lambda_2 = 2$, and $\lambda_3 = 1$) has no dominant eigenvalue.

EXAMPLE 1 Finding a Dominant Eigenvalue

Find the dominant eigenvalue and corresponding eigenvectors of the matrix

$$A = \begin{bmatrix} 2 & -12 \\ 1 & -5 \end{bmatrix}.$$

Solution From Example 4 of Section 7.1 you know that the characteristic polynomial of A is $\lambda^2 + 3\lambda + 2 = (\lambda + 1)(\lambda + 2)$. So the eigenvalues of A are $\lambda_1 = -1$ and $\lambda_2 = -2$, of which the dominant one is $\lambda_2 = -2$. From the same example you know that the dominant eigenvectors of A (those corresponding to $\lambda_2 = -2$) are of the form

$$\mathbf{x} = t \begin{bmatrix} 3 \\ 1 \end{bmatrix}, \quad t \neq 0.$$

The Power Method

Like the Jacobi and Gauss-Seidel methods, the power method for approximating eigenvalues is iterative. First assume that the matrix A has a dominant eigenvalue with corresponding dominant eigenvectors. Then choose an initial approximation \mathbf{x}_0 of one of the dominant eigenvectors of A . This initial approximation must be a *nonzero* vector in R^n . Finally, form the sequence given by

$$\begin{aligned} \mathbf{x}_1 &= A\mathbf{x}_0 \\ \mathbf{x}_2 &= A\mathbf{x}_1 = A(A\mathbf{x}_0) = A^2\mathbf{x}_0 \\ \mathbf{x}_3 &= A\mathbf{x}_2 = A(A^2\mathbf{x}_0) = A^3\mathbf{x}_0 \\ &\vdots \\ \mathbf{x}_k &= A\mathbf{x}_{k-1} = A(A^{k-1}\mathbf{x}_0) = A^k\mathbf{x}_0. \end{aligned}$$

For large powers of k , and by properly scaling this sequence, you will see that you obtain a good approximation of the dominant eigenvector of A . This procedure is illustrated in Example 2.

EXAMPLE 2 Approximating a Dominant Eigenvector by the Power Method

Complete six iterations of the power method to approximate a dominant eigenvector of

$$A = \begin{bmatrix} 2 & -12 \\ 1 & -5 \end{bmatrix}.$$

Solution Begin with an initial nonzero approximation of

$$\mathbf{x}_0 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}.$$

Then obtain the following approximations.

| Iteration | “Scaled” Approximation |
|---|--|
| $\mathbf{x}_1 = A\mathbf{x}_0 = \begin{bmatrix} 2 & -12 \\ 1 & -5 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} -10 \\ -4 \end{bmatrix}$ | → $-4 \begin{bmatrix} 2.50 \\ 1.00 \end{bmatrix}$ |
| $\mathbf{x}_2 = A\mathbf{x}_1 = \begin{bmatrix} 2 & -12 \\ 1 & -5 \end{bmatrix} \begin{bmatrix} -10 \\ -4 \end{bmatrix} = \begin{bmatrix} 28 \\ 10 \end{bmatrix}$ | → $10 \begin{bmatrix} 2.80 \\ 1.00 \end{bmatrix}$ |
| $\mathbf{x}_3 = A\mathbf{x}_2 = \begin{bmatrix} 2 & -12 \\ 1 & -5 \end{bmatrix} \begin{bmatrix} 28 \\ 10 \end{bmatrix} = \begin{bmatrix} -64 \\ -22 \end{bmatrix}$ | → $-22 \begin{bmatrix} 2.91 \\ 1.00 \end{bmatrix}$ |
| $\mathbf{x}_4 = A\mathbf{x}_3 = \begin{bmatrix} 2 & -12 \\ 1 & -5 \end{bmatrix} \begin{bmatrix} -64 \\ -22 \end{bmatrix} = \begin{bmatrix} 136 \\ 46 \end{bmatrix}$ | → $46 \begin{bmatrix} 2.96 \\ 1.00 \end{bmatrix}$ |
| $\mathbf{x}_5 = A\mathbf{x}_4 = \begin{bmatrix} 2 & -12 \\ 1 & -5 \end{bmatrix} \begin{bmatrix} 136 \\ 46 \end{bmatrix} = \begin{bmatrix} -280 \\ -94 \end{bmatrix}$ | → $-94 \begin{bmatrix} 2.98 \\ 1.00 \end{bmatrix}$ |
| $\mathbf{x}_6 = A\mathbf{x}_5 = \begin{bmatrix} 2 & -12 \\ 1 & -5 \end{bmatrix} \begin{bmatrix} -280 \\ -94 \end{bmatrix} = \begin{bmatrix} 568 \\ 190 \end{bmatrix}$ | → $190 \begin{bmatrix} 2.99 \\ 1.00 \end{bmatrix}$ |

Note that the approximations in Example 2 appear to be approaching scalar multiples of

$$\begin{bmatrix} 3 \\ 1 \end{bmatrix},$$

which you know from Example 1 is a dominant eigenvector of the matrix

$$A = \begin{bmatrix} 2 & -12 \\ 1 & -5 \end{bmatrix}.$$

In Example 2 the power method was used to approximate a dominant eigenvector of the matrix A . In that example you already knew that the dominant eigenvalue of A was $\lambda = -2$. For the sake of demonstration, however, assume that you do not know the dominant eigenvalue of A . The following theorem provides a formula for determining the eigenvalue corresponding to a given eigenvector. This theorem is credited to the English physicist John William Rayleigh (1842–1919).

Theorem 10.2

Determining an Eigenvalue from an Eigenvector

If \mathbf{x} is an eigenvector of a matrix A , then its corresponding eigenvalue is given by

$$\lambda = \frac{A\mathbf{x} \cdot \mathbf{x}}{\mathbf{x} \cdot \mathbf{x}}.$$

This quotient is called the **Rayleigh quotient**.

Proof Because \mathbf{x} is an eigenvector of A , you know that $A\mathbf{x} = \lambda\mathbf{x}$ and can write

$$\frac{A\mathbf{x} \cdot \mathbf{x}}{\mathbf{x} \cdot \mathbf{x}} = \frac{\lambda\mathbf{x} \cdot \mathbf{x}}{\mathbf{x} \cdot \mathbf{x}} = \frac{\lambda(\mathbf{x} \cdot \mathbf{x})}{\mathbf{x} \cdot \mathbf{x}} = \lambda.$$

In cases for which the power method generates a good approximation of a dominant eigenvector, the Rayleigh quotient provides a correspondingly good approximation of the dominant eigenvalue. The use of the Rayleigh quotient is demonstrated in Example 3.

EXAMPLE 3 *Approximating a Dominant Eigenvalue*

Use the result of Example 2 to approximate the dominant eigenvalue of the matrix

$$A = \begin{bmatrix} 2 & -12 \\ 1 & -5 \end{bmatrix}.$$

Solution After the sixth iteration of the power method in Example 2, obtained

$$\mathbf{x}_6 = \begin{bmatrix} 568 \\ 190 \end{bmatrix} \approx 190 \begin{bmatrix} 2.99 \\ 1.00 \end{bmatrix}.$$

With $\mathbf{x} = (2.99, 1)$ as the approximation of a dominant eigenvector of A , use the Rayleigh quotient to obtain an approximation of the dominant eigenvalue of A . First compute the product $A\mathbf{x}$.

$$A\mathbf{x} = \begin{bmatrix} 2 & -12 \\ 1 & -5 \end{bmatrix} \begin{bmatrix} 2.99 \\ 1.00 \end{bmatrix} = \begin{bmatrix} -6.02 \\ -2.01 \end{bmatrix}$$

Then, because

$$A\mathbf{x} \cdot \mathbf{x} = (-6.02)(2.99) + (-2.01)(1) \approx -20.0$$

and

$$\mathbf{x} \cdot \mathbf{x} = (2.99)(2.99) + (1)(1) \approx 9.94,$$

you can compute the Rayleigh quotient to be

$$\lambda = \frac{A\mathbf{x} \cdot \mathbf{x}}{\mathbf{x} \cdot \mathbf{x}} \approx \frac{-20.0}{9.94} \approx -2.01$$

which is a good approximation of the dominant eigenvalue $\lambda = -2$.

From Example 2 you can see that the power method tends to produce approximations with large entries. In practice it is best to “scale down” each approximation before proceeding to the next iteration. One way to accomplish this **scaling** is to determine the component of $A\mathbf{x}_i$ that has the largest absolute value and multiply the vector $A\mathbf{x}_i$ by the reciprocal of this component. The resulting vector will then have components whose absolute values are less than or equal to 1. (Other scaling techniques are possible. For examples, see Exercises 27 and 28.)

EXAMPLE 4 *The Power Method with Scaling*

Calculate seven iterations of the power method with *scaling* to approximate a dominant eigenvector of the matrix

$$A = \begin{bmatrix} 1 & 2 & 0 \\ -2 & 1 & 2 \\ 1 & 3 & 1 \end{bmatrix}.$$

Use $\mathbf{x}_0 = (1, 1, 1)$ as the initial approximation.

Solution One iteration of the power method produces

$$A\mathbf{x}_0 = \begin{bmatrix} 1 & 2 & 0 \\ -2 & 1 & 2 \\ 1 & 3 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 3 \\ 1 \\ 5 \end{bmatrix},$$

and by scaling you obtain the approximation

$$\mathbf{x}_1 = \frac{1}{5} \begin{bmatrix} 3 \\ 1 \\ 5 \end{bmatrix} = \begin{bmatrix} 0.60 \\ 0.20 \\ 1.00 \end{bmatrix}.$$

A second iteration yields

$$A\mathbf{x}_1 = \begin{bmatrix} 1 & 2 & 0 \\ -2 & 1 & 2 \\ 1 & 3 & 1 \end{bmatrix} \begin{bmatrix} 0.60 \\ 0.20 \\ 1.00 \end{bmatrix} = \begin{bmatrix} 1.00 \\ 1.00 \\ 2.20 \end{bmatrix}$$

and

$$\mathbf{x}_2 = \frac{1}{2.20} \begin{bmatrix} 1.00 \\ 1.00 \\ 2.20 \end{bmatrix} = \begin{bmatrix} 0.45 \\ 0.45 \\ 1.00 \end{bmatrix}.$$

Continuing this process, you obtain the sequence of approximations shown in Table 10.6.

TABLE 10.6

| \mathbf{x}_0 | \mathbf{x}_1 | \mathbf{x}_2 | \mathbf{x}_3 | \mathbf{x}_4 | \mathbf{x}_5 | \mathbf{x}_6 | \mathbf{x}_7 |
|--|--|--|--|--|--|--|--|
| $\begin{bmatrix} 1.00 \\ 1.00 \\ 1.00 \end{bmatrix}$ | $\begin{bmatrix} 0.60 \\ 0.20 \\ 1.00 \end{bmatrix}$ | $\begin{bmatrix} 0.45 \\ 0.45 \\ 1.00 \end{bmatrix}$ | $\begin{bmatrix} 0.48 \\ 0.55 \\ 1.00 \end{bmatrix}$ | $\begin{bmatrix} 0.51 \\ 0.51 \\ 1.00 \end{bmatrix}$ | $\begin{bmatrix} 0.50 \\ 0.49 \\ 1.00 \end{bmatrix}$ | $\begin{bmatrix} 0.50 \\ 0.50 \\ 1.00 \end{bmatrix}$ | $\begin{bmatrix} 0.50 \\ 0.50 \\ 1.00 \end{bmatrix}$ |

From Table 10.6 you can approximate a dominant eigenvector of A to be

$$\mathbf{x} = \begin{bmatrix} 0.50 \\ 0.50 \\ 1.00 \end{bmatrix}.$$

Using the Rayleigh quotient, you can approximate the dominant eigenvalue of A to be $\lambda = 3$. (For this example you can check that the approximations of \mathbf{x} and λ are exact.)

REMARK: Note that the *scaling factors* used to obtain the vectors in Table 10.6,

| \mathbf{x}_1 | \mathbf{x}_2 | \mathbf{x}_3 | \mathbf{x}_4 | \mathbf{x}_5 | \mathbf{x}_6 | \mathbf{x}_7 |
|----------------|----------------|----------------|----------------|----------------|----------------|----------------|
| ↓ | ↓ | ↓ | ↓ | ↓ | ↓ | ↓ |
| 5.00 | 2.20 | 2.82 | 3.13 | 3.02 | 2.99 | 3.00, |

are approaching the dominant eigenvalue $\lambda = 3$.

In Example 4 the power method with scaling converges to a dominant eigenvector. The following theorem states that a sufficient condition for convergence of the power method is that the matrix A be diagonalizable (and have a dominant eigenvalue).

Theorem 10.3

Convergence of the Power Method

If A is an $n \times n$ diagonalizable matrix with a dominant eigenvalue, then there exists a nonzero vector \mathbf{x}_0 such that the sequence of vectors given by

$$A\mathbf{x}_0, A^2\mathbf{x}_0, A^3\mathbf{x}_0, A^4\mathbf{x}_0, \dots, A^k\mathbf{x}_0, \dots$$

approaches a multiple of the dominant eigenvector of A .

Proof Because A is diagonalizable, you know from Theorem 7.5 that it has n linearly independent eigenvectors $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n$ with corresponding eigenvalues of $\lambda_1, \lambda_2, \dots, \lambda_n$. Assume that these eigenvalues are ordered so that λ_1 is the dominant eigenvalue (with a corresponding eigenvector of \mathbf{x}_1). Because the n eigenvectors $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n$ are linearly independent, they must form a basis for R^n . For the initial approximation \mathbf{x}_0 , choose a nonzero vector such that the linear combination

$$\mathbf{x}_0 = c_1\mathbf{x}_1 + c_2\mathbf{x}_2 + \cdots + c_n\mathbf{x}_n$$

has nonzero leading coefficients. (If $c_1 = 0$, the power method may not converge, and a different \mathbf{x}_0 must be used as the initial approximation. See Exercises 21 and 22.) Now, multiplying both sides of this equation by A produces

$$\begin{aligned} A\mathbf{x}_0 &= A(c_1\mathbf{x}_1 + c_2\mathbf{x}_2 + \cdots + c_n\mathbf{x}_n) \\ &= c_1(A\mathbf{x}_1) + c_2(A\mathbf{x}_2) + \cdots + c_n(A\mathbf{x}_n) \\ &= c_1(\lambda_1\mathbf{x}_1) + c_2(\lambda_2\mathbf{x}_2) + \cdots + c_n(\lambda_n\mathbf{x}_n). \end{aligned}$$

Repeated multiplication of both sides of this equation by A produces

$$A^k\mathbf{x}_0 = c_1(\lambda_1^k\mathbf{x}_1) + c_2(\lambda_2^k\mathbf{x}_2) + \cdots + c_n(\lambda_n^k\mathbf{x}_n),$$

which implies that

$$A^k\mathbf{x}_0 = \lambda_1^k \left[c_1\mathbf{x}_1 + c_2 \left(\frac{\lambda_2}{\lambda_1} \right)^k \mathbf{x}_2 + \cdots + c_n \left(\frac{\lambda_n}{\lambda_1} \right)^k \mathbf{x}_n \right].$$

Now, from the original assumption that λ_1 is larger in absolute value than the other eigenvalues it follows that each of the fractions

$$\frac{\lambda_2}{\lambda_1}, \frac{\lambda_3}{\lambda_1}, \dots, \frac{\lambda_n}{\lambda_1}$$

is less than 1 in absolute value. So each of the factors

$$\left(\frac{\lambda_2}{\lambda_1} \right)^k, \left(\frac{\lambda_3}{\lambda_1} \right)^k, \dots, \left(\frac{\lambda_n}{\lambda_1} \right)^k$$

must approach 0 as k approaches infinity. This implies that the approximation

$$A^k\mathbf{x}_0 \approx \lambda_1^k c_1 \mathbf{x}_1, \quad c_1 \neq 0$$

improves as k increases. Because \mathbf{x}_1 is a dominant eigenvector, it follows that any scalar multiple of \mathbf{x}_1 is also a dominant eigenvector, so showing that $A^k\mathbf{x}_0$ approaches a multiple of the dominant eigenvector of A .

The proof of Theorem 10.3 provides some insight into the rate of convergence of the power method. That is, if the eigenvalues of A are ordered so that

$$|\lambda_1| > |\lambda_2| \geq |\lambda_3| \geq \dots \geq |\lambda_n|,$$

then the power method will converge quickly if $|\lambda_2|/|\lambda_1|$ is small, and slowly if $|\lambda_2|/|\lambda_1|$ is close to 1. This principle is illustrated in Example 5.

EXAMPLE 5 *The Rate of Convergence of the Power Method*

(a) The matrix

$$A = \begin{bmatrix} 4 & 5 \\ 6 & 5 \end{bmatrix}$$

has eigenvalues of $\lambda_1 = 10$ and $\lambda_2 = -1$. So the ratio $|\lambda_2|/|\lambda_1|$ is 0.1. For this matrix, only four iterations are required to obtain successive approximations that agree when rounded to three significant digits. (See Table 10.7.)

TABLE 10.7

| \mathbf{x}_0 | \mathbf{x}_1 | \mathbf{x}_2 | \mathbf{x}_3 | \mathbf{x}_4 |
|--|--|--|--|--|
| $\begin{bmatrix} 1.000 \\ 1.000 \end{bmatrix}$ | $\begin{bmatrix} 0.818 \\ 1.000 \end{bmatrix}$ | $\begin{bmatrix} 0.835 \\ 1.000 \end{bmatrix}$ | $\begin{bmatrix} 0.833 \\ 1.000 \end{bmatrix}$ | $\begin{bmatrix} 0.833 \\ 1.000 \end{bmatrix}$ |

(b) The matrix

$$A = \begin{bmatrix} -4 & 10 \\ 7 & 5 \end{bmatrix}$$

has eigenvalues of $\lambda_1 = 10$ and $\lambda_2 = -9$. For this matrix, the ratio $|\lambda_2|/|\lambda_1|$ is 0.9, and the power method does not produce successive approximations that agree to three significant digits until sixty-eight iterations have been performed, as shown in Table 10.8.

TABLE 10.8

| \mathbf{x}_0 | \mathbf{x}_1 | \mathbf{x}_2 | | \mathbf{x}_{66} | \mathbf{x}_{67} | \mathbf{x}_{68} |
|--|--|--|---------------------|--|--|--|
| $\begin{bmatrix} 1.000 \\ 1.000 \end{bmatrix}$ | $\begin{bmatrix} 0.500 \\ 1.000 \end{bmatrix}$ | $\begin{bmatrix} 0.941 \\ 1.000 \end{bmatrix}$ | $\cdot \cdot \cdot$ | $\begin{bmatrix} 0.715 \\ 1.000 \end{bmatrix}$ | $\begin{bmatrix} 0.714 \\ 1.000 \end{bmatrix}$ | $\begin{bmatrix} 0.714 \\ 1.000 \end{bmatrix}$ |

In this section you have seen the use of the power method to approximate the *dominant* eigenvalue of a matrix. This method can be modified to approximate other eigenvalues through use of a procedure called **deflation**. Moreover, the power method is only one of several techniques that can be used to approximate the eigenvalues of a matrix. Another popular method is called the **QR algorithm**.

This is the method used in most computer programs and calculators for finding eigenvalues and eigenvectors. The algorithm uses the *QR*-factorization of the matrix, as presented in Chapter 5. Discussions of the deflation method and the *QR* algorithm can be found in most texts on numerical methods.

SECTION 10.3 EXERCISES

In Exercises 1–6, use the techniques presented in Chapter 7 to find the eigenvalues of the matrix A . If A has a dominant eigenvalue, find a corresponding dominant eigenvector.

$$1. A = \begin{bmatrix} 2 & 1 \\ 0 & -4 \end{bmatrix} \quad 2. A = \begin{bmatrix} -3 & 0 \\ 1 & 3 \end{bmatrix}$$

$$3. A = \begin{bmatrix} 1 & -5 \\ -3 & -1 \end{bmatrix} \quad 4. A = \begin{bmatrix} 4 & -5 \\ 2 & -3 \end{bmatrix}$$

$$5. A = \begin{bmatrix} 2 & 3 & 1 \\ 0 & -1 & 2 \\ 0 & 0 & 3 \end{bmatrix} \quad 6. A = \begin{bmatrix} -5 & 0 & 0 \\ 3 & 7 & 0 \\ 4 & -2 & 3 \end{bmatrix}$$

In Exercises 7–10, use the Rayleigh quotient to compute the eigenvalue λ of A corresponding to the eigenvector \mathbf{x} .

$$7. A = \begin{bmatrix} 4 & -5 \\ 2 & -3 \end{bmatrix}, \mathbf{x} = \begin{bmatrix} 5 \\ 2 \end{bmatrix} \quad 8. A = \begin{bmatrix} 2 & 3 \\ 1 & 4 \end{bmatrix}, \mathbf{x} = \begin{bmatrix} -3 \\ 1 \end{bmatrix}$$

$$9. A = \begin{bmatrix} 1 & 2 & -2 \\ -2 & 5 & -2 \\ -6 & 6 & -3 \end{bmatrix}, \mathbf{x} = \begin{bmatrix} 1 \\ 1 \\ 3 \end{bmatrix}$$

$$10. A = \begin{bmatrix} 3 & 2 & -3 \\ -3 & -4 & 9 \\ -1 & -2 & 5 \end{bmatrix}, \mathbf{x} = \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix}$$

In Exercises 11–14, use the power method with scaling to approximate a dominant eigenvector of the matrix A . Start with $\mathbf{x}_0 = (1, 1)$ and calculate five iterations. Then use \mathbf{x}_5 to approximate the dominant eigenvalue of A .

$$11. A = \begin{bmatrix} 2 & 1 \\ 0 & -7 \end{bmatrix} \quad 12. A = \begin{bmatrix} -1 & 0 \\ 1 & 6 \end{bmatrix}$$

$$13. A = \begin{bmatrix} 1 & -4 \\ -2 & 8 \end{bmatrix} \quad 14. A = \begin{bmatrix} 6 & -3 \\ -2 & 1 \end{bmatrix}$$

In Exercises 15–18, use the power method with scaling to approximate a dominant eigenvector of the matrix A . Start with $\mathbf{x}_0 = (1, 1, 1)$ and calculate four iterations. Then use \mathbf{x}_4 to approximate the dominant eigenvalue of A .

$$15. A = \begin{bmatrix} 3 & 0 & 0 \\ 1 & -1 & 0 \\ 0 & 2 & 8 \end{bmatrix} \quad 16. A = \begin{bmatrix} 1 & 2 & 0 \\ 0 & -7 & 1 \\ 0 & 0 & 0 \end{bmatrix}$$

$$17. A = \begin{bmatrix} -1 & -6 & 0 \\ 2 & 7 & 0 \\ 1 & 2 & -1 \end{bmatrix} \quad 18. A = \begin{bmatrix} 0 & 6 & 0 \\ 0 & -4 & 0 \\ 2 & 1 & 1 \end{bmatrix}$$

In Exercises 19 and 20, the matrix A does not have a dominant eigenvalue. Apply the power method with scaling, starting with $\mathbf{x}_0 = (1, 1, 1)$, and observe the results of the first four iterations.

$$19. A = \begin{bmatrix} 1 & 1 & 0 \\ 3 & -1 & 0 \\ 0 & 0 & -2 \end{bmatrix} \quad 20. A = \begin{bmatrix} 1 & 2 & -2 \\ -2 & 5 & -2 \\ -6 & 6 & -3 \end{bmatrix}$$

21. **Writing** (a) Find the eigenvalues and corresponding eigenvectors of

$$A = \begin{bmatrix} 3 & -1 \\ -2 & 4 \end{bmatrix}.$$

(b) Calculate two iterations of the power method with scaling, starting with $\mathbf{x}_0 = (1, 1)$.

(c) Explain why the method does not seem to converge to a dominant eigenvector.

22. **Writing** Repeat Exercise 21 using $\mathbf{x}_0 = (1, 1, 1)$, for the matrix

$$A = \begin{bmatrix} -3 & 0 & 2 \\ 0 & -1 & 0 \\ 0 & 1 & -2 \end{bmatrix}.$$

23. The matrix

$$A = \begin{bmatrix} 2 & -12 \\ 1 & -5 \end{bmatrix}$$

has a dominant eigenvalue of $\lambda = -2$. Observe that $A\mathbf{x} = \lambda\mathbf{x}$ implies that

$$A^{-1}\mathbf{x} = \frac{1}{\lambda}\mathbf{x}.$$

Apply five iterations of the power method (with scaling) on A^{-1} to compute the eigenvalue of A with the smallest magnitude.

24. Repeat Exercise 23 for the matrix

$$A = \begin{bmatrix} 2 & 3 & 1 \\ 0 & -1 & 2 \\ 0 & 0 & 3 \end{bmatrix}.$$

-  25. (a) Compute the eigenvalues of


$$A = \begin{bmatrix} 2 & 1 \\ 1 & 2 \end{bmatrix} \text{ and } B = \begin{bmatrix} 2 & 3 \\ 1 & 4 \end{bmatrix}.$$

- (b) Apply four iterations of the power method with scaling to each matrix in part (a), starting with $\mathbf{x}_0 = (-1, 2)$.
 (c) Compute the ratios λ_2/λ_1 for A and B . For which do you expect faster convergence?

26. Use the proof of Theorem 10.3 to show that

$$A(A^k \mathbf{x}_0) \approx \lambda_1(A^k \mathbf{x}_0)$$

for large values of k . That is, show that the scale factors obtained in the power method approach the dominant eigenvalue.

-  In Exercises 27 and 28, apply four iterations of the power method (with scaling) to approximate the dominant eigenvalue of the matrix. After each iteration, scale the approximation by dividing by its length so that the resulting approximation will be a unit vector.

27. $A = \begin{bmatrix} 5 & 6 \\ 4 & 3 \end{bmatrix}$

28. $A = \begin{bmatrix} 7 & -4 & 2 \\ 16 & -9 & 6 \\ 8 & -4 & 5 \end{bmatrix}$